Serial No. 10/604,022 Filed: June 23, 2003

Page 2

## Listing of Claims

 (Currently Amended) A process for the solid phase synthesis of peptides, which comprises:

(a) transferring solid phase resin between a resin source external to a single microwave transparent vessel and the microwave transparent vessel;

- (a)(b) deprotecting a first amino acid linked to a the solid phase resin by removing protective chemical groups from said the first acid;
- (b) (c) activating chemical groups on a second amino acid to prepare the second amino acid for coupling with the first amino acid;
- (e)(d) coupling the activated second amino acid to the deprotected first amino acid to form a peptide from the first and second amino acids;
- (d)(c) accelerating at least the deprotecting and coupling steps by applying microwave energy during the deprotecting, and coupling steps; and
- (e) (f) successively deprotecting, activating, and coupling a plurality of amino acids into a peptide in a the single microwave transparent vessel without removing the peptide from the single vessel between cycles.
- (Original) A process according to Claim 1 comprising cleaving the peptide from the solid phase resin while applying microwave energy to accelerate the cleaving step.

## 3. (Canceled)

4. (Previously Presented) A peptide synthesis process according to Claim 1, comprising successively deprotecting, activating, and coupling a plurality of amino acids into a peptide in the single vessel without removing the peptide from the solid phase resin between cycles.

Serial No. 10/604,022 Filed: June 23, 2003

Page 3

- 5. (Original) A peptide synthesis process according to Claim 1, comprising proactively cooling the vessel and its contents during the application of microwave energy to thereby prevent undesired degradation of the peptide or acids by limiting heat accumulation that would otherwise result from the application of the microwave energy.
- (Original) A peptide synthesis process according to Claim I, wherein the deprotecting step comprises deprotecting the alpha-amino group of the amino acid.
- (Original) A peptide synthesis process according to Claim 1, further comprising deprotecting side chains on the amino acids of the peptide under microwave irradiation.
- (Original) A peptide synthesis process according to Claim 1, wherein the activating step comprises activating the alpha-carboxyl group of the second amino acid.
- 9. (Original) A peptide synthesis process according to Claim 1, wherein the step of applying the microwave energy comprises limiting the application of microwave energy (spiking) to relatively short time intervals to thereby prevent undesired degradation of the peptide or acids by limiting heat accumulation that could be encouraged by the continuous application of the microwave energy.
- 10. (Original) A peptide synthesis process according to Claim 1 further comprising activating and coupling using a in situ method and a composition selected from the group consisting of phosphorium activators, uronium activators, HATU, HBTU, PyBOP, PyAOP, and HOBT
- 11. (Original) A peptide syntheses process according to Claim 1 comprising monitoring the temperature of the vessel and moderating the applied power based upon the monitored temperature.

Serial No. 10/604,022 Filed: June 23, 2003

Page 4

12. (Original) A peptide synthesis process according to Claim 1 comprising moderating the applied power based on the status of the reaction.

13-82 (Cancelled).

THE REMAINDER OF THIS PAGE IS INTENTIONALLY LEFT BLANK